

Neutrino Oscillations Induced by Two-loop Radiative Effects

Teruyuki Kitabayashi^{a *} and Masaki Yasue^{b †‡}

^a Accelerator Engineering Center,
Mitsubishi Electric System & Service Engineering Co.Ltd.,
2-8-8 Umezono, Tsukuba, Ibaraki 305-0045, Japan.

^b Department of Natural Science,
School of Marine Science and Technology, Tokai University,
3-20-1 Orido, Shimizu, Shizuoka 424-8610, Japan
and
Department of Physics, Tokai University,
1117 Kitakaname, Hiratsuka, Kanagawa 259-1291, Japan.

TOKAI-HEP/TH-0008, October, 2000

Abstract

Phenomena of neutrino oscillations are discussed on the basis of two-loop radiative neutrino mechanism. Neutrino mixings are experimentally suggested to be maximal in both atmospheric and solar neutrino oscillations. By using $L_e - L_\mu - L_\tau$ ($\equiv L'$)-conservation, which, however, only ensures the maximal solar neutrino mixing, we find that two-loop radiative mechanism dynamically generates the maximal atmospheric neutrino mixing and that the estimate of $\Delta m_\odot^2/\Delta m_{atm}^2 \sim \epsilon m_e/m_\tau$ explains $\Delta m_\odot^2/\Delta m_{atm}^2 \ll 1$ because of $m_e/m_\tau \ll 1$, where ϵ measures the breaking of the L' -conservation. Together with $\Delta m_{atm}^2 \approx 3 \times 10^{-3} \text{ eV}^2$, this estimate yields $\Delta m_\odot^2 \sim 10^{-7} \text{ eV}^2$ for $\epsilon \sim 0.1$, which corresponds to the LOW solution to the solar neutrino problem. Neutrino mass scale is given by $(16\pi^2)^{-2} m_e m_\tau / M$ ($M \sim 1 \text{ TeV}$), which is of order 0.01 eV.

1 Introduction

Neutrino oscillations have been long recognized if neutrinos are massive particles [1]. Such oscillations in fact have been recently confirmed by the Super-Kamiokande collaboration [2] and have also been observed for solar neutrinos produced inside the Sun [3]. The recent report from the K2K collaboration [4] has further shown that the atmospheric neutrino oscillations are characterized by $\Delta m_{atm}^2 \approx 3 \times 10^{-3} \text{ eV}^2$, which implies $\sim 5.5 \times 10^{-2} \text{ eV}$ as neutrino masses. This tiny mass scale for neutrinos can be generated by radiative mechanisms, where the smallness originates from the smallness of radiative effects [7, 8]. Radiative mechanisms uses $L=2$ interactions given by $\nu_L^{[i} \ell_L^{j]}$ for one-loop radiative effects [7, 9, 10, 11] and by additional $\ell_R^{\{i} \ell_R^{j\}}$ for two-loop radiative effects [8, 12], where i and j denote three families ($i, j = 1, 2, 3$).

At the one-loop level, Zee [7] has presented the mechanism that utilizes a new standard Higgs scalar called ϕ' in addition to the standard Higgs scalar, ϕ , both of which are $SU(2)_L$ -doublets, and another

*E-mail:teruyuki@post.kek.jp

†E-mail:yasue@keyaki.cc.u-tokai.ac.jp

‡A talk given at *Post Summer Institute 2000 on Neutrino Physics*, August 21-24, 2000, Fuji-Yoshida, Japan.




Figure 1: Radiatively generated Majorana neutrino masses: (a) one-loop diagram, (b) two-loop diagram.

singly charged scalar called h^+ , which is an $SU(2)_L$ -singlet, with the coupling of $f_{[ij]}\nu_L^{[i}\ell_L^{j]}h^+$. The Fermi statistics forces $\nu_L^{[i}\ell_L^{j]}$ to be antisymmetrized with respect to the family indices. After the spontaneous breakdown of $SU(2)_L \times U(1)_Y$, an interaction of $\phi\phi'h^+$ yields the possible mixing of h^+ with ϕ^+ characterized by the scale of μ , which finally induces Majorana neutrino masses. Again, the Fermi statistics forces $\phi\phi'$ to be antisymmetrized with respect to the $SU(2)_L$ -indices. Depicted in Figure 1(a) is the diagram for generating Majorana neutrino masses. The order-of-magnitude estimate gives the one-loop neutrino mass, m_ν^{1-loop} , for $\nu_i-\nu_j$ to be:

$$m_\nu^{1-loop} \sim f_{[ij]} \frac{m_{\ell_j}^2}{16\pi^2 M^2} \mu, \quad (1)$$

for $\langle 0|\phi^0|0\rangle \sim \langle 0|\phi'^0|0\rangle$, where M stands for the scale of the model, presumably of order 1 TeV. The factor of $16\pi^2$ in the denominator is specific to one-loop radiative corrections. This estimate turns out to be

$$m_\nu^{1-loop} \sim 2 \times 10^3 f_{[i\tau]} \left(\frac{\mu}{100 \text{ GeV}} \right) \text{ eV}, \quad (2)$$

for $m_{\ell_j} = m_\tau$ ($j=\tau$). To obtain $m_\nu \sim 0.1$ eV, we require that

$$f_{[i\tau]} \sim 5 \times 10^{-5}, \quad (3)$$

for $\mu \sim 100$ GeV. Therefore, to get tiny neutrino masses of order 0.1 eV, one has to give excessive suppression to the lepton-number violating $\nu\ell$ -coupling.

At the two-loop level, additional suppression arises. In addition to h^+ , a doubly charged k^{++} -scalar is required to realize the mechanism of the Zee-Babu type [8] and k^{++} couples to a right-handed charged lepton pair via $\ell_R^{\{i}\ell_R^{j\}}k^{++}$ with coupling strength of $f_{\{ij\}}$. Using a possible coupling of this new k^{++} with h^+ via $h^+h^+k^{++\dagger}$, we can find interactions corresponding to Figure 1(b). The order-of-magnitude estimate gives the two-loop neutrino mass, m_ν^{2-loop} , for $\nu_i-\nu_k$ to be:

$$m_\nu^{2-loop} \sim f_{[ij]} f_{\{jj'\}} f_{[kj']} \frac{m_{\ell_j} m_{\ell_{j'}}}{(16\pi^2)^2 M^2} \mu. \quad (4)$$

The factor of $(16\pi^2)^2$ in the denominator is specific to two-loop radiative corrections. This estimate turns out to yield

$$m_\nu^{2-loop} \sim 10 f_{[i\tau]} f_{[\tau\tau]} f_{[j\tau]} \left(\frac{\mu}{100 \text{ GeV}} \right) \text{ eV}, \quad (5)$$

for $m_{\ell j, \ell j'} = m_\tau$ ($j, j' = \tau$). To obtain $m_\nu \sim 0.1$ eV, thanks to the extra loop-factor of $16\pi^2$, we only require that

$$f_{[i\tau]} \sim 0.1, \quad (6)$$

for $f_{[\tau\tau]} \sim 1$ and $\mu \sim 100$ GeV. Therefore, the two-loop radiative neutrino masses can be of order of 0.1 eV without excessive suppression for relevant couplings [13].

2 Bimaximal Mixing

The observed pattern of neutrino oscillations is consistent with the pattern arising from the requirement of the conservation of the new quantum number $L_e - L_\mu - L_\tau (\equiv L')$ [14]. The $U(1)_{L'}$ symmetry based on the L' -conservation can be used to describe the bimaximal mixing scheme for neutrino oscillations [15, 16]. However, the L' -conservation itself only ensures the maximal solar neutrino mixing but does not determine the atmospheric neutrino mixing angle. In fact, in the one-loop radiative mechanism, fine-tuning of lepton-number violating couplings is necessary to yield bimaximal mixing for atmospheric neutrino oscillations.

In the one-loop radiative mechanism, we have known the form of the neutrino mass matrix, which is given by

$$M_\nu \propto \left(\begin{array}{ccc} 0 & f_{[e\mu]} m_\mu^2 & f_{[e\tau]} m_\tau^2 \\ & 0 & f_{[\mu\tau]} m_\tau^2 \\ & & 0 \end{array} \right) \Big|_{sym} \Rightarrow \left(\begin{array}{ccc} 0 & \sim 1 & \sim 1 \\ & 0 & \varepsilon (\ll 1) \\ & & 0 \end{array} \right) m, \quad (7)$$

where m stands for the neutrino mass scale. The bimaximal mixing is realized if the couplings satisfy

$$f_{e\mu} m_\mu^2 = f_{[e\tau]} m_\tau^2 \Rightarrow f_{[e\mu]} \gg f_{[e\tau]} (\gg f_{[\mu\tau]} \approx 0), \quad (8)$$

indicating the fine-tuning of the couplings f 's. This fine-tuning is referred to as “inverse hierarchy in the couplings”, namely, $f_{[e\mu]} \gg f_{[e\tau]}$ [17]. The L' -conservation gives $f_{[\mu\tau]} = 0$. Its tiny breaking effect characterized by the parameter, ε , produces tiny solar neutrino oscillations.

On the other hand, in the two-loop radiative mechanism, we will find the mass matrix [12] given by

$$M_\nu \propto \left(\begin{array}{ccc} 0 & f_{[e\tau]} f_{[e\mu]} m_e m_\tau & f_{[e\tau]} f_{[e\tau]} m_e m_\tau \\ & f_{[e\mu]} f_{[e\mu]} m_e^2 & f_{[e\mu]} f_{[e\tau]} m_e^2 \\ & & f_{[e\tau]} f_{[e\tau]} m_e^2 \end{array} \right) \Big|_{sym} \Rightarrow \left(\begin{array}{ccc} 0 & \sim 1 & \sim 1 \\ & \varepsilon & \varepsilon' \\ & & \varepsilon'' \end{array} \right) m. \quad (9)$$

The bimaximal structure is reproduced if

$$f_{[e\tau]} f_{[e\mu]} m_e m_\tau = f_{[e\tau]} f_{[e\tau]} m_e m_\tau \Rightarrow f_{[e\mu]} = f_{[e\tau]}. \quad (10)$$

Therefore, no hierarchy in the couplings is necessary. The breaking of the L' -conservation gives the suppressed entries, $\varepsilon, \varepsilon', \varepsilon''$, proportional to m_e^2 . Therefore, we observe that

$$\Delta m_\odot^2 / \Delta m_{atm}^2 \propto m_e / m_\tau, \quad (11)$$

which dynamically guarantees $\Delta m_{atm}^2 \gg \Delta m_\odot^2$ because of $m_\tau \gg m_e$.

In radiative mechanisms, the hierarchy of $\Delta m_{atm}^2 \gg \Delta m_\odot^2$ can also be ascribed to the generic smallness of two-loop radiative effects over one-loop radiative effects [18]. Therefore, we have in hands two dynamical reasons for $\Delta m_{atm}^2 \gg \Delta m_\odot^2$:

$$\frac{\Delta m_\odot^2}{\Delta m_{atm}^2} \ll 1 \text{ because } \left\{ \begin{array}{l} 2\text{-loop}/1\text{-loop} \ll 1 \\ m_e/m_\tau \ll 1 \end{array} \right. . \quad (12)$$

Table 1: L and L' quantum numbers.

Fields	$(\nu_{eL}, e_L^-), e_R^-$	$(\nu_{iL}, \ell_{iL}^-), \ell_{iR}^- _{i=\mu, \tau}$	ϕ	h^+	k^{++}	k'^{++}
L	1	1	0	-2	-2	-2
L'	1	-1	0	0	0	-2

3 Two-loop Radiative Neutrino Masses

Interactions that we introduce can be classified by the ordinary lepton number (L) and L' -number of particles, which are listed in the Table 1. The new ingredients that are not contained in the standard model are the $SU(2)_L$ -singlet scalars, h^+ and k^{++} . We have further employed an additional k^{++} to be denoted by k'^{++} in order to import the L' -breaking. The L - and L' -quantum number of k'^{++} is also listed in Table 1. Extra L - and L' -conserving Yukawa interactions are given by

$$\left\{ \begin{array}{l} f_{[ej]} \left(\nu_{eL} \ell_L^j - \nu_L^j e_L^- \right) h^+, \\ f_{\{ej\}} e_R^- \ell_R^j k^{++}, \\ \frac{1}{2} f_{\{ee\}} e_R^- e_R^- k'^{++}. \end{array} \right. \quad (13)$$

An L -breaking but L' -conserving interaction is specified by

$$\mu_0 h^+ h^+ k^{++\dagger}, \quad (14)$$

where μ_0 represents a mass scale. An L' -breaking interaction is activated by k'^{++} via

$$\mu_b h^+ h^+ k'^{++\dagger}, \quad (15)$$

where μ_b represents a breaking scale of the L' -conservation.

Yukawa interactions, then, take the form of ¹

$$\begin{aligned} -\mathcal{L}_Y &= \sum_{i=e, \mu, \tau} f_\phi^i \overline{\psi_L^i} \phi \ell_R^i + \sum_{i=\mu, \tau} \left(f_{[ei]} \overline{(\psi_L^e)^c} \psi_L^i h^+ + f_{\{ei\}} \overline{(e_R)^c} \ell_R^i k^{++} \right) \\ &+ \frac{1}{2} f_{\{ee\}} \overline{(e_R)^c} e_R k'^{++} + (\text{h.c.}), \end{aligned} \quad (16)$$

and Higgs interactions are described by self-Hermitian terms composed of $\varphi \varphi^\dagger$ ($\varphi = \phi, h^+, k^{++}, k'^{++}$) and by the non-self-Hermitian terms in

$$V_0 = \mu_0 h^+ h^+ k^{++\dagger} + (\text{h.c.}). \quad (17)$$

This coupling softly breaks the L -conservation but preserves the L' -conservation. To account for solar neutrino oscillations, the breaking of the L' -conservation should be included and is assumed to be furnished by

$$V_b = \mu_b h^+ h^+ k'^{++\dagger} + (\text{h.c.}). \quad (18)$$

Neutrino masses are generated by interactions corresponding to the diagrams depicted in Figure 2(a,b). The resulting Majorana neutrino mass matrix is given by

$$M_\nu = \begin{pmatrix} 0 & m_{e\mu} & m_{e\tau} \\ m_{e\mu} & \delta_{\mu\mu} & \delta_{\mu\tau} \\ m_{e\tau} & \delta_{\mu\tau} & \delta_{\tau\tau} \end{pmatrix}. \quad (19)$$

¹The corresponding expression of Eq.(2) in Ref.[19] should read this equation.




Figure 2: Radiatively generated Majorana neutrino masses: (a) L' -conserving two-loop diagram, (b) L' -breaking two-loop diagram.

Here, the bimaximal structure is controlled by

$$m_{ei} \approx -2f_{[e\tau]}f_{[ei]}f_{\{\tau e\}} \frac{m_\tau m_e}{m_k^2} \mu_0 \left[\frac{1}{16\pi^2} \ln \left(\frac{m_k^2}{m_h^2} \right) \right]^2 \quad (i = \mu, \tau), \quad (20)$$

where the product of m_e and m_τ appears. This is because the exchanged leptons are e and τ as can be seen from Figure 2(a). Tiny splitting is induced by

$$\delta_{ij} \approx -f_{[ei]}f_{[ej]}f_{\{ee\}} \frac{m_e m_e}{m_{k'}^2} \mu_b \left[\frac{1}{16\pi^2} \ln \left(\frac{m_{k'}^2}{m_h^2} \right) \right]^2, \quad (21)$$

where m_e^2 appears because the exchanged leptons are both e and e as can be seen from Figure 2(b). These expressions, Eqs.(20) and (21), are subject to the approximation of $m_{k,k'}^2 \gg (\text{other mass squared})$. The detailed derivation of Eqs.(20) and (21) can be found in the Appendix of Ref.[19]. Oscillations are described by these mass parameters:

$$\Delta m_{atm}^2 = m_{e\mu}^2 + m_{e\tau}^2 (\equiv m_\nu^2), \quad \Delta m_\odot^2 = 4m_\nu \delta m, \quad (22)$$

where

$$\delta m = \frac{1}{2} |\delta_{\mu\mu} \cos^2 \theta_\nu + 2\delta_{\mu\tau} \cos \theta_\nu \sin \theta_\nu + \delta_{\tau\tau} \sin^2 \theta_\nu| \quad (23)$$

with

$$\cos \theta_\nu = m_{e\mu}/m_\nu, \quad \sin \theta_\nu = m_{e\tau}/m_\nu. \quad (24)$$

It is thus found that (nearly) bimaximal mixing is reproduced by requiring

$$f_{[e\mu]} \approx f_{[e\tau]}, \quad (25)$$

yielding $\sin 2\theta_\nu \approx 1$. Tiny mass-splitting $\Delta m_{atm}^2 \gg \Delta m_\odot^2$ is ensured by the mass-hierarchy:

$$m_\tau \gg m_e. \quad (26)$$

As a result, we obtain an estimate of the ratio:

$$\frac{\Delta m_\odot^2}{\Delta m_{atm}^2} \sim \frac{\mu_b}{\mu_0} \frac{m_e}{m_\tau} \frac{m_k^2}{m_{k'}^2}. \quad (27)$$




Figure 3: (a) $\mu^- \rightarrow e^- e^- e^+$, (b) $\mu^- \rightarrow e^- \gamma$.




Figure 4: (a) $\tau^- \rightarrow \mu^- e^- e^+$, (b) $\tau^- \rightarrow \mu^- \gamma$.

From this estimate, we find that

$$\begin{aligned}
 \Delta m_\odot^2 &\sim 3 \times 10^{-4} \frac{\mu_b}{\mu_0} \Delta m_{atm}^2 \quad (m_k^2 \sim m_{k'}^2) \\
 \Rightarrow \Delta m_\odot^2 &\sim 3 \times 10^{-5} \Delta m_{atm}^2 \quad (\mu_b \sim \mu_0/10) \\
 \Rightarrow \Delta m_\odot^2 &\sim 10^{-7} \text{ eV}^2 \quad (\Delta m_{atm}^2 \sim 3 \times 10^{-3} \text{ eV}^2).
 \end{aligned} \tag{28}$$

The resulting Δm_\odot^2 corresponds to the allowed region for the LOW solution to the solar neutrino problem. Since k^{++} and k'^{++} couple to the charged lepton pairs, these scalars produce extra contributions on the well-established low-energy phenomenology. In particular, we should consider effects from $\mu^- \rightarrow e^- \gamma$, $e^- e^- e^+$, $e^- e^- \rightarrow e^- e^-$ and $\nu_\mu e^- \rightarrow \nu_\mu e^-$. The relevant constraints on the parameters associated with the scalars of h^+ , k^{++} and k'^{++} are, thus, given by ²

1. $\mu^- \rightarrow e^- e^- e^+$ in Figure 3(a) and $\mu^- \rightarrow e^- \gamma$ in Figure 3(b) [20] (forbidden by the L' -conservation), yielding

$$\frac{\xi f_{e\mu} f_{ee}}{\tilde{m}_k^2} < \begin{cases} 1.2 \times 10^{-10} \text{ GeV}^{-2} \text{ from } B(\mu^- \rightarrow e^- e^- e^+) < 10^{-12} \text{ [21]} \\ 2.4 \times 10^{-8} \text{ GeV}^{-2} \text{ from } B(\mu^- \rightarrow e^- \gamma) < 1.2 \times 10^{-11} \text{ [21]} \end{cases}, \tag{29}$$

²The constraints of Eqs.(12) and (13) in Ref.[19] should, respectively, be replaced by the corresponding bounds in the items 1, 3 and 4. Namely, $f_{\{11,12\}}$ should read $f_{\{11,12\}}/2$ in Ref.[19].




Figure 5: (a) $e^-e^- \rightarrow e^-e^-$, (b) $\nu_\mu e^- \rightarrow \nu_\mu e^-$.

where $\bar{m}_k \sim m_k \sim m_{k'}$ and ξ estimated to be

$$\xi \sim \frac{1}{16\pi^2} \frac{\mu_b \mu_0}{\bar{m}_k^2} (\ll 1) \quad (30)$$

reads the suppression due to the approximate L' -conservation,

2. $\tau^- \rightarrow \mu^- e^- e^+$ in Figure 4(a) and $\tau^- \rightarrow \mu^- \gamma$ in Figure 4(b) (allowed by the L' -conservation), yielding

$$\begin{aligned} \left| \frac{f_{\{e\tau\}} f_{\{e\mu\}}}{\bar{m}_k^2} \right| &< \begin{cases} 2.1 \times 10^{-7} \text{ GeV}^{-2} \text{ from } B(\tau^- \rightarrow \mu^- e^- e^+) < 1.7 \times 10^{-6} [21] \\ 4.2 \times 10^{-6} \text{ GeV}^{-2} \text{ from } B(\tau^- \rightarrow \mu^- \gamma) < 1.1 \times 10^{-6} [21] \end{cases}, \\ \left| \frac{f_{[e\tau]} f_{[e\mu]}}{m_h^2} \right| &< 4.2 \times 10^{-6} \text{ GeV}^{-2} \text{ from } B(\tau^- \rightarrow \mu^- \gamma), \end{aligned} \quad (31)$$

3. $e^-e^- \rightarrow e^-e^-$ [22] in Figure 5(a), yielding

$$\left| \frac{f_{\{ee\}}}{m_{k'}} \right|^2 < 4.8 \times 10^{-5} \text{ GeV}^{-2}, \quad (32)$$

4. $\nu_\mu e^- \rightarrow \nu_\mu e^-$ [23] in Figure 5(b), yielding

$$\left| \frac{f_{[e\mu]}}{m_h} \right|^2 < 1.7 \times 10^{-6} \text{ GeV}^{-2}. \quad (33)$$

It should be noted that the leading contribution of h^+ to $\mu^- \rightarrow e^- \gamma$, which gives the most stringent constraint on h^+ , is forbidden by the $U(1)_{L'}$ -invariant coupling structure.

Typical parameter values are so chosen to satisfy these constraints:

$$\left. \begin{aligned} &f_{[e\mu]} = f_{[e\tau]} \approx 2e \\ &f_{\{ee\}} = f_{\{e\tau\}} \approx e \end{aligned} \right\} \text{ to suppress higher - order effects,}$$

$$\left. \begin{aligned} &m_h \approx 350 \text{ GeV} \\ &m_k = m_{k'} \approx 2 \text{ TeV} \\ &\mu_0 \approx 1.5 \text{ TeV} \\ &\mu_b \approx \mu_0/10 \end{aligned} \right\} \text{ to suppress exotic contributions.} \quad (34)$$

We obtain the following numerical values:

$$\begin{cases} \Delta m_{atm}^2 \approx 2.4 \times 10^{-3} \text{ eV}^2, \\ \Delta m_{\odot}^2 \approx 10^{-7} \text{ eV}^2. \end{cases} \quad (35)$$

Therefore, we in fact successfully explain phenomena of atmospheric and solar neutrino oscillations characterized by $\Delta m_{atm}^2 \approx 2.4 \times 10^{-3} \text{ eV}^2$ and $\Delta m_{\odot}^2 \approx 10^{-7} \text{ eV}^2$.

4 Summary

We have discussed how neutrino oscillations arise from two loop-radiative mechanism, which exhibits

1. bimaximal mixing due to the $L_e - L_\mu - L_\tau$ conservation via the coupling of $e^- \tau^- k^{++}$,
2. dynamically induced tiny mass-splitting for solar neutrino oscillations due to the smallness of m_e via $e^- e^- k'^{++}$.

The interactions required to generate two-loop Majorana neutrino masses are specified by

$$\begin{cases} f_{[ei]} (\nu_{eL} \ell_L^i - \nu_L^i e_L^-) h^+ \\ f_{\{ei\}} e_R^- \ell_R^i k^{++} \\ \frac{1}{2} f_{\{ee\}} e_R^- e_R^- k'^{++} \end{cases} \quad (i = \mu, \tau) \oplus \begin{cases} \mu_0 h^+ h^+ k^{++\dagger} \\ \mu_b h^+ h^+ k'^{++\dagger} \end{cases} \quad (36)$$

The resulting mass scale for neutrino masses is determined by

$$\frac{m_\tau m_e}{(16\pi^2)^2 m_k^2} \mu_0 \sim \frac{m_\tau m_e}{(16\pi^2)^2 m_k} \sim 10^{-2} \text{ eV}. \quad (37)$$

Thus, to obtain the neutrino mass of order of 0.01 eV is a natural consequence without fine-tuning of coupling parameters. And the hierarchy of $\Delta m_{atm}^2 \gg \Delta m_{\odot}^2$ is expressed by the estimate

$$\Delta m_{\odot}^2 \sim \frac{\mu_b}{\mu_0} \frac{m_e}{m_\tau} \frac{m_k^2}{m_{k'}^2} \Delta m_{atm}^2, \quad (38)$$

which ensures $\Delta m_{atm}^2 \gg \Delta m_{\odot}^2$ because of $m_\tau \gg m_e$.³ This estimation yields the LOW solution to the solar neutrino problem.

It should be finally noted that

- since the L' -conservation forbids primary flavor-changing processes involving e^- , the coupling strengths of h^+ and k^{++} to leptons are not severely constrained and can be as large as $\mathcal{O}(e)$,
- characteristic signatures of h^+ include

$$B(h^+ \rightarrow e^+ \cancel{E}_T) \approx 2B(h^+ \rightarrow \mu^+ \cancel{E}_T) \approx 2B(h^+ \rightarrow \tau^+ \cancel{E}_T) \quad (39)$$

since $f_{[e\mu]} \approx f_{[e\tau]}$, which should be compared with [24]

$$B(h^+ \rightarrow e^+ E_T) \approx B(h^+ \rightarrow \mu^+ \cancel{E}_T) \gg B(h^+ \rightarrow \tau^+ E_T) \quad (40)$$

in the one-loop radiative mechanism with $f_{[e\mu]} \gg f_{[e\tau]} \gg f_{[\mu\tau]}$ [17].

Acknowledgments

We are grateful to M. Matsuda for his suggestion on possible constraints from τ -decays. The work of M.Y. is supported by the Grant-in-Aid for Scientific Research No 12047223 from the Ministry of Education, Science, Sports and Culture, Japan.

³One should be aware of higher-order contributions found by Lavoura in Ref.[12]. The (1,1)-entry of M_ν , which vanishes up to the two-loop level, is induced by the four-loop diagram shown in Figure 6. The contributions are at most characterized by $\delta \sim (16\pi^2)^{-1} \xi m_\tau^2/m_k^2$, which should be compared with $m_e^2/m_{k'}^2$. Our parameter-setting in Eq.(34) gives $\delta \sim 2m_e^2/m_k^2$, which turns out to be $\mathcal{O}(m_e^2/m_{k'}^2)$. Therefore, our estimate of Eq.(38) remains valid to predict Δm_{\odot}^2 from Δm_{atm}^2 .




Figure 6: Four loop-diagram for ν_e - ν_e .

References

- [1] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870. See also B. Pontecorvo, JETP (USSR) **34** (1958) 247; Zh. Eksp. Teor. Piz. **53** (1967) 1717; V. Gribov and B. Pontecorvo, Phys. Lett. **28B** (1969) 493.
- [2] SuperKamiokande Collaboration, Y. Fukuda *et al.*, Phys. Rev. Lett. **81** (1998) 1562; Phys. Lett. B **433** (1998) 9 and **436** (1998) 33. See also K. Scholberg, hep-ex/9905016 (May, 1999). For recent analysis, see N. Forego, M.C. Gonzalez-Garcia and J.W.F. Valle, JHEP **7** (2000) 006; Nucl. Phys. B **580** (2000) 58.
- [3] J. N. Bahcall, P. I. Krastev and A.Yu.Smirnov, Phys. Rev. D **58** (1998) 096016; *ibid.* **60** (1999) 093001; J. N. Bahcall, hep-ph/0002018 (Feb, 2000); M. C. Gonzalez-Garcia, P. C. de Holanda, C. Pena-Garay and J. C. W. Valle, Nucl. Phys. B **573** (2000) 3; G.L. Fogli, E. Lisi, D. Montanino and A. Palazzo, Phys. Rev. D **62** (2000) 013002,
- [4] M. Sakuda, Talk given at *the 30th Int. Conf. on High Energy Physics (ICHEP2000)*, 27 Jul. - 2 Aug., Osaka, Japan. See also, K2K Collaboration, T.Ishida, hep-ex/0008047 (Aug 2000).
- [5] Y. Takeuchi, Talk given at *the 30th Int. Conf. on High Energy Physics (ICHEP2000)*, 27 Jul. - 2 Aug., Osaka, Japan. See also, J. Ellis, hep-ph/0008334 (Aug, 2000).
- [6] T. Yanagida, in *Proceedings of the Workshop on Unified Theories and Baryon Number in the Universe* edited by A. Sawada and A. Sugamoto (KEK Report No.79-18, Tsukuba, 1979), p.95; Prog. Theor. Phys. **64** (1980) 1103; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity* edited by P. van Nieuwenhuizen and D.Z. Freedmann (North-Holland, Amsterdam 1979), p.315; R.N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980) 912.
- [7] A. Zee, Phys. Lett. **93B** (1980) 389; **161B** (1985) 141; L. Wolfenstein, Nucl. Phys. **B175** (1980) 93.
- [8] A. Zee, Nucl. Phys. **B264** (1986) 99; K. S. Babu, Phys. Lett. B **203** (1988) 132; D. Chang, W.-Y.Keung and P.B. Pal, Phys. Rev. Lett. **61** (1988) 2420. See also, J. Schechter and J.W.F. Valle, Phys. Rev. D **25** (1982) 2951.
- [9] S. P. Petcov, Phys. Lett. **115B** (1982) 401; K. S. Babu and V. S. Mathur, Phys. Lett. B **196** (1987) 218; J. Liu, Phys. Lett. B **216** (1989) 367; D. Chang and W.-Y. Keung, Phys. Rev. D **39** (1989) 1386; W. Grimus and H. Neufeld, Phys. Lett. B **237** (1990) 521; B.K. Pal, Phys. Rev. D **44** (1991) 2261; W. Grimus and G. Nardulli, Phys. Lett. B **271** (1991) 161; J.T. Polonium, A. Yu. Smirnov and J.W.F. Valle, Phys. Lett. B **286** (1992) 321; A. Yu. Smirnov and Z. Tao, Nucl. Phys. B **426** (1994) 415.

- [10] A.Yu. Smirnov and M. Tanimoto, Phys. Rev. D **55** (1997) 1665; N. Gaur, A. Ghosal, E. Ma and P. Roy, Phys. Rev D **58** (1998) 071301; Y. Okamoto and M. Yasuè, Prog. Theor. Phys. **101** (1999) 1119; Phys. Lett. B **466** (1999) 267; G.C. McLaughlin and J.N. Ng, Phys. Lett. B **455** (1999) 224; J.E. Kim and J.S. Lee, hep-ph/9907452 (July, 1999); N. Haba, M. Matsuda and M. Tanimoto, Phys. Lett. B **478** (2000) 351; C.-K. Chula, X.-G. He and W.-Y.P. Hwang, Phys. Lett. B **479** (2000) 224; K. Cheung and O.C.W. Kong, Phys. Rev. D **61** (2000) 113012.
- [11] C. Jarlskog, M. Matsuda, S. Skadhauge and M. Tanimoto, Phys. Lett. B **449** (1999) 240; P.H. Frampton and S.L. Glashow, Phys. Lett. B **461** (1999) 95; A.S. Joshipura and S.D. Rindani, Phys. Lett. B **464** (1999) 239; D. Chang and A. Zee, Phys. Rev. D **61** (2000) 071303.
- [12] L. Lavoura, Phys. Rev. D **62** (2000) 093011; T. Kitabayashi and M. Yasuè, Phys. Lett. B **490** (2000) 236.
- [13] See for example, R.N. Mohapatra, hep-ph/9910365 (Oct., 1999).
- [14] R. Barbieri, L. J. Hall, D. Smith, N. J. Weiner and A. Strumia, JHEP **12** (1998) 017. For earlier attempts of using such modified lepton numbers, see for example, S. T. Petcov, Phys. Lett. **110B** (1982) 245; C.N. Leung and S. T. Petcov, Phys. Lett. **125B** (1983) 461; A. Zee, in Ref. [8].
- [15] D.V. Ahluwalia, Mod. Phys. Lett. A **13** (1998) 2249; V. Barger, P. Pakvasa, T.J. Weiler and K. Whisnant, Phys. Lett. B **437** (1998) 107; A. Baltz, A.S. Goldhaber and M. Goldhaber, Phys. Rev. Lett. **81** (1998) 5730; Y. Nomura and T. Yanagida, Phys. Rev. D **59** (1999) 017303; M. Jezabek and Y. Sumino, Phys. Lett. B **440** (1998) 327; R.N. Mohapatra and S. Nussinov, Phys. Lett. B **441** (1998) 299; Q. Shafi and Z. Tavartkiladze, Phys. Lett. B **451** (1999) 129; Phys. Lett. B **482** (2000) 145; I. Stancu and D.V. Ahluwalia, Phys. Lett. B **460** (1999) 431; H. Georgi and S.L. Glashow, Phys. Rev. D **61** (2000) 097301; R. N. Mohapatra, A. Pérez-Lorenzana and C.A. de S. Pires, Phys. Lett. B **474** (2000) 355.
- [16] H. Fritzsch and Z.Z. Xing, Phys. Lett. B **372** (1996) 265; *ibid.* **440** (1998) 313; M. Fukugida, M. Tanimoto and T. Yanagida, Phys. Rev. D **57** (1998) 4429; M. Tanimoto, Phys. Rev. D. **59** (1999) 017304.
- [17] C. Jarlskog, M. Matsuda, S. Skadhauge and M. Tanimoto, in Ref.[11].
- [18] A.S. Joshipura and S.D. Rindani, Phys. Lett. B **464** (1999) 239; D. Chang and A. Zee, Phys. Rev. D **61** (2000) 071303; T. Kitabayashi and M. Yasuè, hep-ph/0006040 (June, 2000); hep-ph/0010087 (Oct., 2000).
- [19] T. Kitabayashi and M. Yasuè, in Ref.[12].
- [20] K. S. Babu, in Ref.[8].
- [21] Particle Data Group, D.E. Groom et al. Euro. Phys. J. C **15** (2000) 1.
- [22] A.S. Joshipura and S.D. Rindani, in Ref.[11].
- [23] A.Yu. Smirnov and M. Tanimoto, in Ref.[10].
- [24] In the Zee model, where $f_{[e\mu]}$ is as small as $\mathcal{O}(10^{-4})$, the h^+ -phenomenology in LEP has been lately focused in the literature. See S. Kanemura, T. Kaai, G.-L. Lin, Y. Okada, J.-J. Tseng and C.-P. Yuan, hep-ph/0010233 (Oct., 2000).